# Author index


Abdolhay, H *see* Alavi, S M H
Acosta, B *see* Méndez, L C
Acosta-Salmón, H & Southgate, P C. Use of a biopsy technique to obtain gonad tissue from the blacklip pearl oyster *Pinctada margaritifera* (L.), 93
Afonso, J M *see* Ginés, R
Agresti, J J *see* Moen, T
Aguirre-Guzmán, G, Ruíz, H M & Ascencio, F. A review of extracellular virulence product of *Vibrio* species important in diseases of cultivated shrimp, 1395
Ahmed, I, Khan, M A & Jafri, A K. Dietary threonine requirement of fingerling Indian major carp, *Cirrhinus mrigala* (Hamilton), 162
Akarasanon, K, Damrongphol, P & Poolsanguan, W. Long-term cryopreservation of spermatophore of the giant freshwater prawn, *Macrobrachium rosenbergii* (de Man), 1415
Alam, M S, Teshima, S-i, Ishikawa, M, Hasegawa, D & Koshio, S. Dietary arginine requirement of juvenile kuruma shrimp *Marsupenaeus japonicus* (Bate), 842
Alavi, S M H, Cosson, J, Karami, M, Abdolhay, H & Amiri, B M. Chemical composition and osmolality of seminal fluid of *Acipenser persicus*: their physiological relationship with sperm motility, 1238
Alcapán, A C *see* Toro, J E
Alexis, M *see* Rigos, G
Alim, M A, Wahab, M A & Milstein, A. Effects of adding different proportions of the small fish punti (*Puntius sophore*) and mola (*Amblypharyngodon mola*) to a polyculture of large carp, 124
Allan, G L & Booth, M A. Effects of extrusion processing on digestibility of peas, lupins, canola meal and soybean meal in silver perch *Bidyanus bidyanus* (Mitchell) diets, 981
Allan, G L & Booth, M A. The effects of dietary digestible protein and digestible energy content on protein retention efficiency of juvenile silver perch *Bidyanus bidyanus* (Mitchell), 970
Allan, G L *see* Booth, M A
Amiri, B M *see* Alavi, S M H
Amirkolaie, A K *see* Schneider, O
Ando, Y & Kobayashi, S. Positional distribution of docosahexaenoic acid in triacyl-*sn*-glycerols of rotifers *Brachionus plicatilis* enriched with fish oil fatty acid ethyl esters, 1391
Andreassen, I *see* Tveiten, H
Aquacop *see* Cuzon, G
Aqutaert, P *see* Harzevili, A S
Arabacı, M, Çağırgan, H & Sarı, M. Induction of ovulation in ornamental common carp (Koi, *Cyprinus carpio* L.) using LHRHa ([D-Ser(tBu)$^{6}$, Pro$^{9}$-NEt]-LHRH) combined with haloperidol and carp pituitary extract, 10
Aralar, E V *see* Santiago, C B
Aranda, D A *see* Patiño-Suárez, V
Arcilla, R P *see* Santiago, C B
Arena, L *see* Cuzon, G
Argüello, A *see* Ginés, R
Arnold, S J *see* Burford, M A
Arredondo-Vega, B O *see* Flores-Vergara, C
Arslan, T & Phelps, R P. Directing gonadal differentiation in bluegill, *Lepomis macrochirus* (Rafinesque), and black crappie, *Pomoxis nigromaculatus* (Lesueur), by periodic estradiol-17β immersions, 397
Arthur, G *see* Fitzsimmons, S
Ascencio, F. *see* Aguirre-Guzmán, G
Attard, M. *see* Harris, J O
Austin, C M *see* Nguyen, T T T
Auwerx, J *see* Harzevili, A S
Avidar, Y. *see* Cnaani, A
Ayyappan, S *see* Das, P C

Babiak, I *see* Rzemieniecki, A
Baghel, D S, Lakra, W S & Rao, G P S. Altered sex ratio in giant fresh water prawn, *Macrobrachium rosenbergii* (de Man) using hormone bioencapsulated live *Artemia* feed, 943
Bai, S C *see* Choi, S-M
Bai, S C *see* Kim, K-W
Bairagi, A, Ghosh, K S, Sen, S K & Ray, A K. Evaluation of the nutritive value of *Leucaena leucocephala* leaf meal, inoculated with fish intestinal bacteria *Bacillus subtilis* and *Bacillus circulans* in formulated diets for rohu, *Labeo rohita* (Hamilton) fingerlings, 436
Bakopoulos, V *see* Poulos, C
Balebona, M C *see* Chabrillón, M
Balogh, K *see* Elbaraasi, H
Barki, A & Karplus, I. Size rank and growth potential in redclaw crayfish (*Cherax quadricarinatus*): are stunted juveniles suitable for grow-out?, 559
Barros, M M *see* Furuya, W M
Barve, S K *see* Shirdhankar, M M
Baylon, J C, Bravo, M E A & Maningo, N C. Ingestion of *Brachionus plicatilis* and *Artemia salina* nauplii by mud crab *Scylla serrata* larvae, 62
Becker, K *see* Schlechtriem, C
Bell, J G *see* Ng, W-K
Benli, A C K & Yildiz, H Y. Blood parameters in Nile tilapia (*Oreochromis niloticus* L.) spontaneously infected with *Edwardsiella tarda*, 1388
Benton, C. *see* DeGraaf, J D
Bercsényi, M *see* Müller, T
Berger, B *see* Lahnsteiner, F
Berger, B *see* Mansour, N
Berlinsky, D L. *see* DeGraaf, J D
Berlinsky, D L *see* Martin-Robichaud, D J
Blier, P U *see* Lamarre, S G
Boonyaratpalin, M *see* Thunjai, T
Booth, M A, Warner-Smith, R J, Allan, G L & Glencross, B D. Effects of dietary astaxanthin source and light manipulation on the skin colour of Australian snapper *Pagrus auratus* (Bloch & Schneider, 1801), 458
Booth, M A *see* Allan, G L
Bordas, M A *see* Chabrillón, M
Borredat, M *see* Vita, R
Boyd, C E *see* Thunjai, T
Bravo, M E A *see* Baylon, J C
Brzuska, E. Artificial spawning of carp (*Cyprinus carpio* L.): differences between the effects of reproduction in females of Hungarian, Polish and French origin treated with carp pituitary homogenate or [D-Tle$^{6}$, ProNHEt$^{9}$] GnRH (*Lecirelin*), 1318
Buckley, L J *see* Peck, M A
Buffa, G *see* Vaccaro, A M
Bunn, S E *see* Burford, M A
Burford, M A, Preston, N P, Minh, T H, Hoa, T T T, Bunn, S E & Fry, V M. Dominant sources of dietary carbon and nitrogen for shrimp reared in extensive rice-shrimp ponds, 194
Burford, M A, Sellars, M J, Arnold, S J, Keys, S J, Crocos, P J & Preston, N P. Contribution of the natural biota associated with substrates to the nutritional requirements of the post-larval shrimp, *Penaeus esculentus* (Haswell), in high-density rearing systems, 508
Burka, J F *see* Westcott, J D
Burridge, L *see* Costello, M J
Burridge, L E, Hamilton, N, Waddy, S L, Haya, K, Mercer, S M, Greenhalgh, R, Tauber, R, Radecki, S V, Crouch, L S, Wislocki, P G & Endris, R G. Acute toxicity of emamectin benzoate (SLICE$^{TM}$) in fish feed to American lobster, *Homarus americanus*, 713

Çağırgan, H *see* Arabacı, M
Cao, L *see* Zheng, D
Carr, J & Whoriskey, F. Sea lice infestation rates on wild and escaped farmed Atlantic salmon (*Salmo salar* L.) entering the Magaguadavic River, New Brunswick, 723
Caruso, G, Maimone, G, Mancuso, M, Modica, A & Genovese, L. Microbiological controls across the productive cycle of *Dicentrarchus labrax* L. and *Sparus aurata* L.: a study from the environment to the final product, 184


© 2004 Blackwell Publishing Ltd

Carvalho, E G *see* Martins da Rocha, R
Catacutan, M R & Pagador, G E. Partial replacement of fishmeal by defatted soybean meal in formulated diets for the mangrove red snapper, *Lutjanus argentimaculatus* (Forsskal 1775), 299
Cernák, I *see* Müller, T
Cerón-Ortiz, A N *see* Flores-Vergara, C
Cesar, A *see* Vita, R
Chabrillón, M, Bordas, M A, Moriñigo, M A & Balebona, M C. Kinetics of adhesion of *Listonella anguillarum* to the mucus of gilt-head seabream, and the implication of surface components, 403
Chadha, N K *see* Khan, M A
Chang, B *see* Costello, M J
Chau, N T H *see* Vromant, N
Chen, Z T, *see* Kang, K H
Cheng, Z J, Hardy, R W & Huige, N J. Apparent digestibility coefficients of nutrients in brewer's and rendered animal by-products for rainbow trout (*Oncorhynchus mykiss* (Walbaum)), 1
Choi, S-M, Wang, X, Park, G-J, Lim, S-R, Kim, K-W, Bai, S C & Shin, I-S. Dietary dehulled soybean meal as a replacement for fish meal in fingerling and growing olive flounder *Paralichthys olivaceus* (Temminck et Schlegel), 410
Choi, S-M *see* Kim, K-W
Choy, E J *see* Jo, Q
Christensen, S M, Macintosh, D J & Phuong, N T, Pond production of the mud crabs *Scylla paramamosain* (Estampador) and *S olivacea* (Herbst) in the Mekong Delta, Vietnam, using two different supplementary diets, 1013
Ciereszko, A *see* Rzemieniecki, A
Civera-Cerecedo, R *see* Cortés-Jacinto, E
Clark, G A *see* Powell, M D
Cnaani, A, Tinman, S, Avidar, Y, Ron, M & Hulata, G, Comparative study of biochemical parameters in response to stress in *Oreochromis aureus*, *O. mossambicus* and two strains of *O. niloticus*, 1434
Cnaani, A *see* Moen, T
Çoban, D *see* Saka, S
Colihueque, N *see* Estay, F J
Coloso, R M *see* Eusebio, P S
Congleton, J L *see* Welker, T L
Cordero-Esquivel, B *see* Flores-Vergara, C
Cordero-Esquivel, B *see* Lora-Vilchis, M C
Cortés-Jacinto, E, Villarreal-Colmenares, H, Civera-Cerecedo, R & Naranjo-Páramo, J, Effect of dietary protein level on the growth and survival of pre-adult freshwater crayfish *Cherax quadricarinatus* (von Martens) in monosex culture, 71
Cosson, J *see* Alavi, S M H
Costello, M J, Burridge, L, Chang, B & Robichaud, L, Sea Lice 2003—Proceedings of the sixth international conference on sea lice biology and control, 711
Coyle, S D, Mengel, G J, Tidwell, J H & Webster, C D, Evaluation of growth, feed utilization, and economics of hybrid tilapia, *Oreochromis niloticus* × *Oreochromis aureus*, fed diets containing different protein sources in combination with distillers dried grains with solubles, 365
Crocco, C *see* Turchini, G M
Crocos, P J *see* Burford, M A
Crouch, L S *see* Burridge, L E
Cruz-Suárez, L-E *see* Tapia-Salazar, M
Csengeri, I *see* Elbaraasi, H
Cui, Y *see* Xue, M
Curnow, J *see* Glencross, B
Cuzon, G, Arena, L, Goguenheim, J M, Goyard, E, & Aquacop, Is it possible to raise, offspring of the 25th generation of *Litopenaeus vannamei* (Boone) and 18th generation *Litopenaeus stylirostris* (Stimpson) in clear water to 40 g?, 1244
Cyrino, J E P *see* Portz, L

Dabrowski, K *see* Lee, K-J
da Conceição, L E C *see* Dias, J
Damrongphol, P, *see* Akarasanon, K
Das, B K *see* Das, P C
Das, P C, Ayyappan, S, Jena, J K & Das, B K, Acute toxicity of ammonia and its sub-lethal effects on selected haematological and enzymatic parameters of mrigal, *Cirrhinus mrigala* (Hamilton), 134
Das, P C, Ayyappan, S, Jena, J K & Das, B K, Effect of sub-lethal nitrite on selected haematological parameters in fingerling *Catla catla* (Hamilton), 874
Davies, E J *see* Peck, M A
de Castro-Mallare, T R *see* Guanzon, N G
De Charleroy, D *see* Harzevili, A S
de Groot, S J, Book review, 1108
de la Parra, I A *see* Hernández, C
De Silva, S S, Gunasekera, R M & Ingram, B A, Performance of intensively farmed Murray cod *Maccullochella peelii peelii* (Mitchell) fed newly formulated vs currently used commercial diets, and a comparison of fillet composition of farmed and wild fish, 1039
Debowski, M *see* Krzemieniewski, M
Decamp, O *see* Nagano, N
Degani, G, Application of RAPD and cytochrome *b* sequences to the study of genetic variations in $F_1$ and $F_2$ generations of two different strains of guppy (*Poecilia reticulata*, Peter 1854), 807
Degani, G *see* Tzchori, I
DeGraaf, J D, King, W, Benton, C & Berlinsky, D L, Production and storage of sperm from the black sea bass *Centropristis striata* L., 1457
Demaine, H *see* Pant, J
Dias, J, Rueda-Jasso, R, Panserat, S, da Conceição, L E C, Gomes, E F & Dinis, M T, Effect of dietary carbohydrate-to-lipid ratios on growth, lipid deposition and metabolic hepatic enzymes in juvenile Senegalese sole (*Solea senegalensis*, Kaup), 1122
Díaz, A C & Fenucci, J L, Effect of prepared diet on the induction of precocious maturation in *Pleoticus muelleri* Bate (Crustacea, Penaeoidea), 1166
Dimitriadis, G J *see* Poulos, C
Dinis, M T *see* Dias, J
Dixon, B A, Shinn, A P & Sommerville, C, Genetic characterization of populations of the ectoparasitic caligid, *Lepeophtheirus salmonis* (Kroyer 1837) using randomly amplified polymorphic DNA, 730
Domagała, J *see* Rzemieniecki, A
Dong, P *see* Qin, J G
Dooremont, I *see* Harzevili, A S
Doupé, R G & Lymbery, A J, Indicators of genetic variation for feed conversion efficiency in black bream, 1305
Dumas, S *see* Peña, R
Duncan, P *see* Erler, D

Eckhoff, K *see* Hemre, G-I
Eding, E H *see* Schneider, O
Edwards, P *see* Pant, J
Egea, Mª A *see* Rondán, M
Eiriksson, H, *see* Jónasson, J P
Eklund, M W, Poysky, F T, Paranjpye, R N, Peterson, M E & Pelroy, G A, Susceptibility of coho salmon, *Oncorhynchus kisutch* (Walbaum), to different toxins of *Clostridium botulinum*, 594
Elbaraasi, H, Mézes, M, Balogh, K, Horváth, L & Csengeri, I, Effects of dietary ascorbic acid/iron ratio on some production traits, lipid peroxide state and amount/activity of the glutathione redox system in African catfish *Clarias gariepinus* (Burchell) fingerlings, 256
El-Gohary, F A *see* El-Shafai, S A
Elisha, R *see* Tzchori, I
Eliyahu, R *see* Tzchori, I
Elnady, M *see* Hassanien, H A
El-Saidy, D M S D & Gaber, M M, Use of cottonseed meal supplemented with iron for detoxification of gossypol as a total replacement of fish meal in Nile tilapia, *Oreochromis niloticus* (L.) diets, 859
El-Shafai, S A, El-Gohary, F A, Verreth, J A J, Schrama, J W & Gijzen, H J, Apparent digestibility coefficient of duckweed (*Lemna minor*), fresh and dry for Nile tilapia (*Oreochromis niloticus* L.), 574
Endris, R G *see* Burridge, L E
Engler, L S *see* Thompson, K R
Erler, D, Pollard, P, Duncan, P & Knibb, W, Treatment of shrimp farm effluent with omnivorous finfish and artificial substrates, 816
Escutia-Zuñiga, S *see* Hernandez-Llamas, A



© 2004 Blackwell Publishing Ltd

Estay, F J, Noriega, R, Ureta, J P, Martín, W & Colihueque, N. Reproductive performance of cultured brown trout (*Salmo trutta* L.) in Chile, 447
Eusebio, P S, Coloso, R M & Mamauag, R E P. Apparent digestibility of selected ingredients in diets for juvenile grouper, *Epinephelus coioides* (Hamilton), 1261
Ezaz, M T, McAndrew, B J & Penman, D J. Spontaneous diploidization of the maternal chromosome set in Nile tilapia (*Oreochromis niloticus* L.) eggs, 271
Ezaz, M T, Sayeed, S, McAndrew, B J & Penman, D J. Use of microsatellite loci and AFLP markers to verify gynogenesis and clonal lines in Nile tilapia *Oreochromis niloticus* L., 1472

Fagbenro, O A. Soybean meal replacement by roquette (*Eruca sativa* Miller) seed meal as protein feedstuff in diets for African Catfish, *Clarias gariepinus* (Burchell 1822), fingerlings, 917
Falk-Petersen, I-B *see* Lamarre, S G
Famula, T R *see* Moen, T
Felton, S M *see* Halver, J E
Fenucci, J L *see* Díaz, A C
Fenucci, J L *see* Gimenez, A V F
Fırat, K *see* Saka, S
Fitzsimmons, S, Saravanan, S, Walden, J & Arthur, G. Effects of delay of onset of feeding and *Artemia* concentrations on the survival and growth of Stage I European lobsters, *Homarus gammarus* (Linnaeus), 605
Flores-Nava, A *see* Martínez-Cárdenas, L
Flores-Vergara, C, Cordero-Esquivel, B, Cerón-Ortiz, A N & Arredondo-Vega, B O. Combined effects of temperature and diet on growth and biochemical composition of the Pacific oyster *Crassostrea gigas* (Thunberg) spat, 1131
Focken, U *see* Schlechtriem, C
Fox, S P *see* Ogilvie, S C
Fry, V M *see* Burford, M A
Furuya, V R B *see* Furuya, W M
Furuya, W M, Pezzato, L E, Barros, M M, Pezzato, A C, Furuya, V R B & Miranda, E C. Use of ideal protein concept for precision formulation of amino acid levels in fish-meal-free diets for juvenile Nile tilapia (*Oreochromis niloticus* L.), 1110

Gabaudan, J *see* Vielma, J
Gaber, M M *see* El-Saidy, D M S D
Gall, G A E *see* Moen, T
García, B *see* Rondán, M
García, G *see* Peña, R
García-Melys, D *see* Winkler, F M
Garduño-Lugo, M, Muñoz-Córdova, G & Olvera-Novoa, M A. Mass selection for red colour in *Oreochromis niloticus* (Linnaeus 1758), 340
Gatlin III, D M *see* Li, P
Genovese, L *see* Caruso, G
Gensic, M, Wissing, P J, Keefe, T R & Mustafa, A. Effects of iodized feed on stress modulation in steelhead trout, *Oncorhynchus mykiss* (Walbaum), 1117
Gettinby, G *see* McKenzie, E
Ghosh, K S *see* Bairagi, A
Gijzen, H J *see* El-Shafai, S A
Gillibrand, P A *see* Penston, M J
Gimenez, A V F, Fenucci, J L & Petriella, A M. The effect of vitamin E on growth, survival and hepatopancreas structure of the Argentine red shrimp *Pleoticus muelleri* Bate (Crustacea, Penaeidea), 1172
Ginés, R, Afonso, J M, Argüello, A, Zamorano, M J & López, J L. The effects of long-day photoperiod on growth, body composition and skin colour in immature gilthead sea bream (*Sparus aurata* L.), 1207
Glencross, B D *see* Booth, M A
Glencross, B, Hawkins, W & Curnow, J. Nutritional assessment of Australian canola meals. I. Evaluation of canola oil extraction method and meal processing conditions on the digestible value of canola meals fed to the red seabream (*Pagrus auratus*, Paulin), 15
Glencross, B, Hawkins, W & Curnow, J. Nutritional assessment of Australian canola meals. II. Evaluation of the influence of the canola oil extraction method on the protein value of canola meals fed to the red seabream (*Pagrus auratus*, Paulin), 25
Glogowski, J *see* Rzemieniecki, A
Goguenheim, J *see* Cuzon, G
Gomes, E F *see* Dias, J
Gomez, C *see* Lee, K-J
Gong, J *see* Liu, S
Gonzal, A C *see* Santiago, C B
Gonzalez-Becerril, A *see* Hernandez-Llamas, A
González-Rodríguez, B *see* Hernández, C
Goyard, E *see* Cuzon, G
Green, T J *see* Harris, J O
Greenhalgh, R *see* Burridge, L E
Grier, H J *see* Tiersch, T R
Grigoriou, P and Richardson, C A. Aspects of the growth of cultured cuttlefish *Sepia officinalis* (Linnaeus 1758), 1141
Grubišić, L *see* Tičina, V
Guanzon Jr, N G, de Castro-Mallare, T R & Lorque, F M. Polyculture of milkfish *Chanos chanos* (Forsskal) and the red seaweed *Gracilariopsis bailinae* (Zhang *et* Xia) in brackish water earthen ponds, 423
Gunasekera, R M *see* De Silva, S S
Guo, X *see* Yang, H
Guo, Z, *see* Lin, H Z
Guz, L *see* Lee, K-J

Halver, J E, Felton, S M & Zbanyszek, R. Carcass selenium loss as an indicator of stress in barge transported chinook salmon (*Oncorhynchus tshawytscha* Walbaum) smolts, 1099
Hamáčkova, J *see* Rzemieniecki, A
Hamilton, N *see* Burridge, L E
Hammell, K L *see* Westcott, J D
Han, D, Xie, S, Lei, W, Zhu, X & Yang, Y. Effect of ration on the growth and energy budget of Chinese longsnout catfish, *Leiocassis longirostris* Günther, 866
Haniffa, M A, Sridhar, S & Nagarajan, M. Induction of triploidy and tetraploidy in stinging catfish, *Heteropneustes fossilis* (Bloch), using heat shock, 937
Harboe, T *see* van der Meeren, T
Hardy, R W *see* Cheng, Z J
Harmon, P *see* Peterson, R H
Harris, A *see* Tapia-Salazar, M
Harris, J O, Powell, M D, Attard, M & Green, T J. Efficacy of chloramine-T as a treatment for amoebic gill disease (AGD) in marine Atlantic salmon (*Salmo salar* L.), 1448
Harvey-Clark, C *see* Roy, P K
Harzevili, A S, Dooremont, I, Vught, I, Auwerx, J, Quataert, P & De Charleroy, D. First feeding of burbot, *Lota lota* (Gadidae, Teleostei) larvae under different temperature and light conditions, 49
Harzevili, A S, Dooremont, I, Vught, I, Auwerx, J, Aqutaert, P & De Charleroy, D. First feeding of burbot, *Lota lota* (Gadidae, Teleostei) larvae under different temperature and light conditions, 419
Hasegawa, D *see* Alam, M S
Hassanien, H A, Elnady, M, Obeida, A & Itriby, H. Genetic diversity of Nile tilapia populations revealed by randomly amplified polymorphic DNA (RAPD), 587
Hattori, M, Sawada, Y, Kurata, M, Yamamoto, S, Kato, K & Kumai, H. Oxygen deficiency during somitogenesis causes centrum defects in red sea bream, *Pagrus major* (Temminck et Schlegel), 850
Hawkins, W *see* Glencross, B
Hay, D W *see* McKibben, M A
Hay, D W *see* Penston, M J
Haya, K *see* Burridge, L E
Heasman, M *see* Liu, W
Hemre, G-I, Karlsen, Ø, Eckhoff, K, Tveit, K, Mangor-Jensen, A & Rosenlund, G. Effect of season, light regime and diet on muscle composition and selected quality parameters in farmed Atlantic cod, *Gadus morhua* L., 683
Hernández-Ceballos, D *see* Peña, R
Hernández, Mª D *see* Rondán, M
Hernández, C, Sarmiento-Pardo, J, González-Rodríguez, B & de la Parra, I A. Replacement of fish meal with co- extruded wet tuna viscera and corn meal in diets for white shrimp (*Litopenaeus vannamei* Boone), 1153
Hernandez-Llamas, A, Gonzalez-Becerril, A, Hernandez-Vazquez, S & Escutia-Zuñiga, S. Bioeconomic analysis of intensive production of the blue shrimp *Litopenaeus stylirostris* (Stimpson), 103

© 2004 Blackwell Publishing Ltd 

Hernandez-Llamas, A *see* Villarreal, H
Hernandez-Vazquez, S *see* Hernandez-Llamas, A
Herrera, A M *see* Liñán-Cabello, M A
Hoa, T T T *see* Burford, M A
Holloway, A C, Keene, J L, Noakes, D G & Moccia, R D. Effects of clove oil and MS-222 on blood hormone profiles in rainbow trout *Oncorhynchus mykiss*, Walbaum, 1025
Horvath, A *see* Lahnsteiner, F
Horváth, L *see* Elbaraasi, H
Hoskonen, P & Pirhonen, J. The effect of clove oil sedation on oxygen consumption of six temperate-zone fish species, 922
Hu, X *see* Liu, S
Huang, C-H & Lin, W-Y. Effects of dietary vitamin E level on growth and tissue lipid peroxidation of soft-shelled turtle, *Pelodiscus sinensis* (Wiegmann), 948
Huige, N J *see* Cheng, Z J
Hulata, G *see* Cnaani, A
Hulata, G *see* Moen, T
Hurvitz, A *see* Tzchori, I
Hussenot, J *see* Kassila, J
Hwang, H *see* Liu, S

Ingram, B A *see* De Silva, S S
Ishikawa, M *see* Alam, M S
Islam, Md. S & Tanaka, M. Optimization of dietary protein requirement for pond-reared mahseer *Tor putitora* Hamilton (Cypriniformes: Cyprinidae), 1270
Itriby, H *see* Hassanien, H A

Jackson, C, Preston, N & Thompson, P J. Intake and discharge nutrient loads at three intensive shrimp farms, 1053
Jackson, S *see* Samocha, T M
Jafri, A K *see* Ahmed, I
Jafri, A K *see* Khan, M A
Jain, K K *see* Venkat, H K
James, M R *see* Ogilvie, S C
Jayasankar, P. Random amplified polymorphic DNA (RAPD) fingerprinting resolves species ambiguity of domesticated clown fish (genus: *Amphiprion*, family: Pomacentridae) from India, 1006
Jee, J-H & Kang, J-C. Effect of phenanthrene on haematological parameters in olive flounder, *Paralichthys olivaceus* (Temminch et Schlegel), 1310
Jena, J K *see* Das, P C
Jiang, X *see* Liu, S
Jiménez-Brinquis, B *see* Vita, R
Jo, Q, Choy, E J, Park, D W & Véron, B. Sterol dynamics of heterotrophic *Tetraselmis suecica* and its nutritional implication in the bivalve aquaculture, 371
Jo, S-G & Marcus, N H. The effect of temperature and duration of incubation on the hatching of diapause eggs of *Centropages hamatus* (Lilljeborg) (Copepoda, Calanoida), 432
Jobling, M. 'Finishing' feeds for carnivorous fish and the fatty acid dilution model, 706
Jobling, M *see* Tveiten, H
Jónasson, J P, Thórarinsdóttir, G G, Eiríksson, H & Marteinsdóttir, G. Temperature tolerance of Iceland scallop, *Chlamys islandica* (O.F. Müller) under controlled experimental conditions, 1405
Jones, E R *see* Samocha, T M
Jover, M *see* Rondán, M

Kang, J-C *see* Jee, J-H
Kang, J-C *see* Kim, S-G
Kang, K H & Kim, J M. The predation of trumpet shell, *Charonia* sp., on eight different marine invertebrate species, 1202
Kang, K H, Kho, K H, Chen, Z T, Kim, J M, Kim, Y H & Zhang, Z F. Cryopreservation of filefish (*Thamnaconus septentrionalis* Gunther, 1877) sperm, 1429
Kang, K H, Shao, M Y, Kho, K H & Zhang, Z F. Short-term storage and cryopreservation of *Urechis unicinctus* (Echiura: Urechidae) sperm, 1195
Karami, M *see* Alavi, S M H
Karlsen, Ø *see* Hemre, G-I
Karplus, I *see* Barki, A
Kassila, J & Hussenot, J. Fractionation of phosphate in marine aquaculture sediments: optimization of the ethylenediaminetetraacetic acid method and comparison with other procedures, 1339
Katavić, I *see* Tičina, V
Kato, K *see* Hattori, M
Kause, A, Ritola, O & Paananen, T. Breeding for improved appearance of large rainbow trout in two production environments, 924
Keefe, T R *see* Gensic, M
Keene, J L *see* Holloway, A C
Keys, S J *see* Burford, M A
Khan, M A, Jafri, A K & Chadha, N K. Growth, reproductive performance, muscle and egg composition in grass carp, *Ctenopharyngodon idella* (Valenciennes), fed hydrilla or formulated diets with varying protein levels, 1277
Khan, M A *see* Ahmed, I
Kho, K H *see* Kang, K H
Kim, J M *see* Kang, K H
Kim, J-W *see* Kim, S-G
Kim, K-W, Wang, X, Choi, S-M, Park, G-J & Bai, S C. Evaluation of optimum dietary protein-to-energy ratio in juvenile olive flounder *Paralichthys olivaceus* (Temminck et Schlegel), 250
Kim, K-W *see* Choi, S-M
Kim, S-G, Kim, J-W & Kang, J-C. Effect of dietary cadmium on growth and haematological parameters of juvenile rockfish, *Sebastes schlegeli* (Hilgendorf), 80
Kim, Y H, *see* Kang, K H
King, T L *see* Silverstein, J T
King, W, *see* DeGraaf, J D
Klesius, P H *see* Shelby, R A
Knibb, W *see* Erler, D
Kobayashi, S *see* Ando, Y
Koshio, S *see* Alam, M S
Kouřil, J *see* Rzemieniecki, A
Krzemieniewski, M, Teodorowicz, M, Debowski, M & Pesta, J. Effect of a constant magnetic field on water quality and rearing of European sheatfish *Silurus glanis* L., larvae, 568
Kucska, B *see* Müller, T
Kumai, H *see* Hattori, M
Kurata, M *see* Hattori, M

Lahnsteiner, F, Berger, B, Horvath, A & Urbanyi, B. Studies on the semen biology and sperm cryopreservation in the sterlet, *Acipenser ruthenus* L., 519
Lahnsteiner, F *see* Mansour, N
Lakra, W S *see* Baghel, D S
Lalitha, K V & Surendran, P K. Bacterial microflora associated with farmed freshwater prawn *Macrobrachium rosenbergii* (de Man) and the aquaculture environment, 629
Lall, S P *see* Roy, P K
Lall, S P *see* Tibbetts, S M
Lamarre, S G, Le François, N R, Falk-Petersen, I-B & Blier, P U. Can digestive and metabolic enzyme activity levels predict growth rate and survival of newly hatched Atlantic wolffish (*Anarhichas lupus* Olafsen)?, 608
Lapolla, A E *see* Peck, M A
Lavkulich, L M *see* Schendel, E K
Lawler, N F *see* O'Connor, W A
Lawrence, A L *see* Samocha, T M
Le François, N R *see* Lamarre, S G
Lee, K-J, Dabrowski, K, Rinchard, J, Gomez, C, Guz, L & Vilchez, C. Supplementation of maca (*Lepidium meyenii*) tuber meal in diets improves growth rate and survival of rainbow trout *Oncorhynchus mykiss* (Walbaum) alevins and juveniles, 215
Legendre, M *see* Nguenga, D
Lei, W *see* Han, D
Levinton, J S *see* Wong, W H
Li, P, Wang, X & Gatlin III, D M. Excessive dietary levamisole suppresses growth performance of hybrid striped bass, *Morone chrysops* × *M. saxatilis*, and elevated levamisole *in vitro* impairs macrophage function, 1380
Li, Z J, *see* Lin, H Z
Lim, S-R *see* Choi, S-M
Lin, H R *see* Wen, H S
Lin, H Z, Guo, Z, Yang, Y, Zheng, W & Li, Z J. Effect of dietary probiotics on apparent digestibility coefficients of nutrients of white shrimp *Litopenaeus vannamei* Boone, 1441
Lin, J *see* Rhyne, A L



© 2004 Blackwell Publishing Ltd

Lin, W-Y *see* Huang, C-H
Liñán-Cabello, M A, Medina-Zendejas, R, Sánchez-Barajas, M & Herrera, A M. Effects of carotenoids and retinol in oocyte maturation of crayfish *Cherax quadrucarinatus*, 905
Ling, N *see* Willis, K J
Liu, S, Jiang, X, Hu, X, Gong, J, Hwang, H & Mai, K. Effects of temperature on non-specific immune parameters in two scallop species: *Argopecten irradians* (Lamarck 1819) and *Chlamys farreri* (Jones & Preston 1904), 678
Liu, S *see* Zheng, D
Liu, W, Heasman, M & Simpson, R. Evaluation of cytochalasin B (CB) treatments for triploidy induction in the blacklip abalone, *Haliotis rubra* (Leach, 1814), 1062
Liu, W, Heasman, M & Simpson, R. Optimization of triploidy induction in the blacklip abalone, *Haliotis rubra* (Leach, 1814), using 6-dimethylaminopurine, 1076
Liufu, Z *see* Zheng, D
Lopez, I M *see* Samocha, T M
López, J L *see* Ginés, R
Lora-Vilchis, M C, Cordero-Esquivel, B & Voltolina, D. Growth of *Artemia franciscana* fed *Isochrysis* sp. and *Chaetoceros muelleri* during its early life stages, 1086
Lorque, F M *see* Guanzon, N G
Luckenbach, J A & Sullivan, C V. Effective GnRHa dose and gamete ratio for reproduction of southern flounder, *Paralichthys lethostigma* (Jordan and Gilbert 1884), 1482
Lupatsch, I *see* Sklan, D
Lymbery, A J *see* Doupé, R G

Macintosh, D J *see* Christensen, S M
Mai, K *see* Liu, S
Mai, K *see* Zheng, D
Maimone, G *see* Caruso, G
Mamauag, R E P *see* Eusebio, P S
Mamuris, Z *see* Moutou, K A
Mancuso, M *see* Caruso, G
Mangor-Jensen, A *see* Hemre, G-I
Maningo, N C *see* Baylon, J C
Mansour, N, Lahnsteiner, F & Berger, B. Characterization of the testicular semen of the African catfish, *Clarias gariepinus* (Burchell, 1822), and its short-term storage, 232
Marín, A *see* Vita, R
Marín-Guirao, L *see* Vita, R
Marcus, N H *see* Jo, S-G
Martín, W *see* Estay, F J
Martínez, F J *see* Rondán, M
Martínez-Cárdenas, L & Flores-Nava, A. Do granular substrates enhance microbiota availability and growth in cultured *Rana catesbeiana* tadpoles?, 616
Marteinsdóttir, G, *see* Jónasson, J P
Martin-Robichaud, D J & Berlinsky, D L. The effects of photothermal manipulation on reproductive development in female haddock *Melanogrammus aeglefinus* L., 465
Martin-Robichaud, D J *see* Peterson, R H
Martins da Rocha, R, Carvalho, E G & Urbinati, E C. Physiological responses associated with capture and crowding stress in matrinxã *Brycon cephalus* (Gunther, 1869), 245
May, B *see* Moen, T
Mazzola, A *see* Vaccaro, A M
McAndrew, B J *see* Ezaz, M T
McCart, K *see* McKenzie, E
McKenzie, E, Gettinby, G, McCart, K & Revie, C W. Time-series models of sea lice *Caligus elongatus* (Nordmann) abundance on Atlantic salmon *Salmo salar* L. in Loch Sunart, Scotland, 764
McKibben, M A & Hay, D W. Distributions of planktonic sea lice larvae *Lepeophtheirus salmonis* in the inter-tidal zone in Loch Torridon, Western Scotland in relation to salmon farm production cycles, 742
McKibben, M A *see* Penston, M J
McVicar, A H. Management actions in relation to the controversy about salmon lice infections in fish farms as a hazard to wild salmonid populations, 751
Medina-Zendejas, R *see* Liñán-Cabello, M A
Méndez, L C, Racotta, I S, Acosta, B & Portillo-Clark, G. Effect of sediment on growth and survival of post-larval *Litopenaeus stylirostris* (Boone, 1931), 652
Mengel, G J *see* Coyle, S D
Mentasti, T *see* Turchini, G M
Mercer, S M *see* Burridge, L E
Mézes, M *see* Elbaraasi, H
Millan, A *see* Villarreal, H
Milley, J E *see* Tibbetts, S M
Milstein, A *see* Alim, M A
Minařík, B *see* Řehulka, J
Minh, T H *see* Burford, M A
Miranda, E C *see* Furuya, W M
Mirto, S *see* Vaccaro, A M
Moav, B *see* Tzchori, I
Moccia, R D *see* Holloway, A C
Modica, A *see* Caruso, G
Moen, T, Agresti, J J, Cnaani, A, Moses, H, Famula, T R, Hulata, G, Gall, G A E & May, B. A genome scan of a four-way tilapia cross supports the existence of a quantitative trait locus for cold tolerance on linkage group 23, 893
Molina-Poveda, C & Morales, M E. Use of a mixture of barley-based fermented grains and wheat gluten as an alternative protein source in practical diets for *Litopenaeus vannamei* (Boone), 1158
Morales, M E *see* Molina-Poveda, C
Moretti, V M *see* Turchini, G M
Moriñigo, M A *see* Chabrillón, M
Morton, S-R *see* Thompson, K R
Moses, H *see* Moen, T
Moutou, K A, Panagiotaki, P & Mamuris, Z. Effects of salinity on digestive protease activity in the euryhaline sparid *Sparus aurata* L.: a preliminary study, 912
Müller, T, Taller, J, Nyitrai, G, Kucska, B, Cernák, I & Bercsényi, M. Hybrid of pikeperch, *Sander lucioperca* L. and Volga perch, *S volgense* (Gmelin), 915
Muñoz-Córdova, G *see* Garduño-Lugo, M
Mustafa, A *see* Gensic, M
Muzinic, L A *see* Thompson, K R
Myers, L J *see* Shelby, R A

Nagano, N & Decamp, O. Ingestion of a ciliated protozoa by first-feeding larval stage of Pacific white shrimp, *Litopenaeus vannamei* (Boone), 516
Nagarajan, M *see* Haniffa, M A
Nakano, T, *see* Shiozaki, K
Nam, C Q *see* Vromant, N
Naranjo-Páramo, J *see* Cortés-Jacinto, E
Nathanailides, C *see* Paschos, I
Neidig, C L *see* Tiersch, T R
Nengas, I *see* Rigos, G
Ng, W-K, Sigholt, T & Bell, J G. The influence of environmental temperature on the apparent nutrient and fatty acid digestibility in Atlantic salmon (*Salmo salar* L.) fed finishing diets containing different blends of fish oil, rapeseed oil and palm oil, 1228
Nguenga, D, Teugels, G G, Legendre, M & Ollevier, F. Effects of storage and incubation temperature on the viability of eggs, embryos and larvae in two strains of an African catfish, *Heterobranchus longifilis* (Siluriformes, Clariidae), 1358
Nguenga, D, Teugels, G G, Legendre, M & Ollevier, F. Influence of tropical seasonal changes on oocyte diameter, responses to hormonal induction and hatching quality in two strains of the catfish, *Heterobranchus longifilis* Val. (Clariidae), 1349
Nguyen, T T T & Austin, C M. Inheritance of molecular markers and sex in the Australian freshwater crayfish, *Cherax destructor* Clark, 1328
Nienhuis, P H. Book review, 1010
Noakes, D G *see* Holloway, A C
Nordstrom, S E *see* Schendel, E K
Noriega, R *see* Estay, F J
Nyitrai, G *see* Müller, T

Obeida, A *see* Hassanien, H A
O'Bryan, L M *see* Peck, M A
O'Connor, W A & Lawler, N F. Reproductive condition of the pearl oyster, *Pinctada imbricata*, Röding, in Port Stephens, New South Wales, Australia, 385
Ogilvie, S C, Fox, S P, Ross, A H, James, M R & Schiel, D R. Growth of cultured mussels (*Perna canaliculus* Gmelin 1791) at a deep-water chlorophyll maximum layer, 1253

© 2004 Blackwell Publishing Ltd

Ojeda, J A *see* Toro, J E
Oliva-Teles, A & Pimentel-Rodrigues, A. Phosphorus requirement of European sea bass (*Dicentrarchus labrax* L.) juveniles, 636
Oliva-Teles, A *see* Pereira, T G
Ollevier, F *see* Nguenga, D
Ollevier, F *see* Vromant, N
Olvera-Novoa, M A *see* Garduño-Lugo, M
Opitz, H M *see* Pietrak, M
Opstad, I & Rust, M B. Effect of recovery salinity on survival of acutely stressed halibut (*Hippoglossus hippoglossus L.*) larvae, 1286

Paananen, T *see* Kause, A
Pagador, G E *see* Catacutan, M R
Palma-Rojas, C *see* Winkler, F M
Panagiotaki, P *see* Moutou, K A
Panserat, S *see* Dias, J
Pant, J, Demaine, H & Edwards, P. Assessment of the aquaculture subsystem in integrated agriculture–aquaculture systems in Northeast Thailand, 289
Paranjpye, R N *see* Eklund, M W
Park, D W *see* Jo, Q
Park, G-J *see* Choi, S-M
Park, G-J *see* Kim, K-W
Paschos, I, Nathanailides, C, Perdikaris, C & Tsoumani, M. Comparison of morphology, growth and survival between *Silurus glanis*, *S. aristotelis* and their hybrid during larval and juvenile stages, 97
Patiño-Suárez, V, Aranda, D A & Zamora, A G. Food ingestion and digestibility of five unicellular algae by 1-day-old *Strombus gigas* larvae, 1149
Peck, M A, Buckley, L J, O'Bryan, L M, Davies, E J & Lapolla, A E. Efficacy of egg surface disinfectants in captive spawning Atlantic cod *Gadus morhua* L. and haddock *Melanogrammus aeglefinus* L., 992
Pelroy, G A *see* Eklund, M W
Peña, R, Dumas, S, Saldivar-Lucio, R, García, G, Trasviña, A & Hernández-Ceballos, D. The effect of light intensity on first feeding of the spotted sand bass *Paralabrax maculatofasciatus* (Steindachner) larvae, 345
Penman, D J *see* Ezaz, M T
Penston, M J, McKibben, M A, Hay, D W & Gillibrand, P A. Observations on open-water densities of sea lice larvae in Loch Shieldaig, Western Scotland, 793
Perdikaris, C *see* Paschos, I
Pereira, T G & Oliva-Teles, A. Evaluation of micronized lupin seed meal as an alternative protein source in diets for gilthead sea bream *Sparus aurata* L. juveniles, 828
Pesta, J *see* Krzemieniewski, M
Peterson, M E *see* Eklund, M W
Peterson, R H, Martin-Robichaud, D J & Harmon, P. Influence of incubation temperature on body movements of Atlantic cod (*Gadus morhua* L.) embryos and on size at hatch, 453
Petriella, A M *see* Gimenez, A V F
Pezzato, A C *see* Furuya, W M
Pezzato, L E *see* Furuya, W M
Phelps, R P *see* Arslan, T
Phuong, N T *see* Christensen, S M
Pietrak, M & Opitz, H M. An evaluation of three potential methods for preventing the spread of larval *Lepeophtheirus salmonis* (Kroyer, 1837), 759
Pimentel-Rodrigues, A *see* Oliva-Teles, A
Pirhonen, J *see* Hoskonen, P
Pollard, P *see* Erler, D
Poolsanguan, W, *see* Akarasanon, K
Portillo-Clark, G *see* Méndez, L C
Portz, L & Cyrino, J E P. Digestibility of nutrients and amino acids of different protein sources in practical diets by largemouth bass *Micropterus salmoides* (Lacepède, 1802), 312
Poulos, C, Bakopoulos, V, Zolota, V & Dimitriadis, G J. Histopathological findings after sea bass (*Dicentrarhus labrax* L.) exposure to extracellular products of *Photobacterium damsela* subsp *piscicida* produced *in vivo*, 931
Powell, M D & Clark, G A. Efficacy and toxicity of oxidative disinfectants for the removal of gill amoebae from the gills of amoebic gill disease affected Atlantic salmon (*Salmo salar* L.) in freshwater, 112
Powell, M D, *see* Harris, J O
Poysky, F T *see* Eklund, M W
Prag, T *see* Sklan, D
Preston, N *see* Jackson, C
Preston, N P *see* Burford, M A
Puzzi, C *see* Turchini, G M

Qin, J G & Dong, P. Acute toxicity of trichlorfon to juvenile yabby *Cherax destructor* (Clark) and selected zooplankton species, 1104
Quataert, P *see* Harzevili, A S

Racotta, I S *see* Méndez, L C
Radecki, S V *see* Burridge, L E
Rao, G P S *see* Baghel, D S
Ray, A K *see* Bairagi, A
Řehulka, J, Minařík, B & Řehulkova, E. Red blood cell indices of rainbow trout *Oncorhynchus mykiss* (Walbaum) in aquaculture, 529
Řehulková, E *see* Řehulka, J
Revie, C W *see* McKenzie, E
Rexroad III, C E *see* Silverstein, J T
Rhyne, A L & Lin, J. Effects of different diets on larval development in a peppermint shrimp (*Lysmata* sp. (Risso)), 1179
Ricci, M *see* Schlechtriem, C
Richardson, C A *see* Grigoriou, P
Ricque-Marie, D *see* Tapia-Salazar, M
Ridha, M T. Observations on the reproductive performance of three mouth-brooding tilapia species in low-salinity underground water, 1031
Rigos, G, Tyrpenou, A E, Nengas, I, Alexis, M & Troisi, G M. The kinetic profile of oxolinic acid in sharpsnout sea bream, *Diplodus puntazzo* (Cetti 1777), 1299
Rinchard, J *see* Lee, K-J
Ritola, O *see* Kause, A
Rivera, M C *see* Villarreal, H
Robichaud, L *see* Costello, M J
Rocha, S *see* Villarreal, H
Ron, M, *see* Cnaani, A
Rondán, M, Hernández, Mª D, Egea, Mª A, García, B, Jover, M, Rueda, F M & Martínez, F J. Effects of fishmeal replacement with soybean meal as protein source, and protein replacement with carbohydrates as an alternative energy source on sharpsnout sea bream, *Diplodus puntazzo*, fatty acid profile, 1220
Rosenlund, G *see* Hemre, G-I
Ross, A H *see* Ogilvie, S C
Roy, P K, Lall, S P & Harvey-Clark, C. A cannulation technique for urine collection from haddock, *Melanogrammus aeglefinus* L., 87
Rueda, F M *see* Rondán, M
Rueda-Jasso, R *see* Dias, J
Ruiz, H M, *see* Aguirre-Guzmán, G
Ruohonen, K *see* Vielma, J
Rust, M B *see* Opstad, I
Rzemieniecki, A, Domagała, J, Glogowski, J, Ciereszko, A, Trzebiatowski, R, Kouřil, J, Hamáčkova, J & Babiak, I. Induced spermiation in 3-year-old sterlet, *Acipenser ruthenus* L., 144

Sahu, N P *see* Venkat, H K
Saka, S, Fırat, K & Çoban, D. Development of the common dentex (*Dentex dentex*) eggs in relation to temperature, 224
Sala, T *see* Turchini, G M
Saldivar-Lucio, R *see* Peña, R
Samocha, T M, Lopez, I M, Jones, E R, Jackson, S & Lawrence, A L. Characterization of intake and effluent waters from intensive and semi-intensive shrimp farms in Texas, 321
Sánchez-Barajas, M *see* Liñán-Cabello, M A
Santiago, C B, Gonzal, A C, Aralar, E V & Arcilla, R P. Effect of stunting of juvenile bighead carp *Aristichthys nobilis* (Richardson) on compensatory growth and reproduction, 836
Sarà, G *see* Vaccaro, A M
Sarı, M *see* Arabacı, M
Saravanan, S *see* Fitzsimmons, S
Sarmiento-Pardo, J *see* Hernández, C
Sato, M, *see* Shiozaki, K



© 2004 Blackwell Publishing Ltd

Sato, N. *see* Shiozaki, K
Sawada, Y *see* Hattori, M
Sayeed, S. *see* Ezaz, M T
Schendel, E K. Nordstrom, S E & Lavkulich, L M. Floc and sediment properties and their environmental distribution from a marine fish farm, 483
Schiel, D R *see* Ogilvie, S C
Schlechtriem, C, Ricci, M, Focken, U & Becker, K. Mass produced nematodes *Panagrellus redivivus* as live food for rearing carp larvae: preliminary results, 547
Schneider, O, Amirkolaie, A K, Vera-Cartas, J, Eding, E H, Schrama, J W & Verreth, J A J. Digestibility, faeces recovery, and related carbon, nitrogen and phosphorus balances of five feed ingredients evaluated as fishmeal alternatives in Nile tilapia, *Oreochromis niloticus* L., 1370
Schrader, K K *see* Shelby, R A
Schrama, J W *see* El-Shafai, S A
Schrama, J W *see* Schneider, O
Sellars, M J *see* Burford, M A
Sen, S K *see* Bairagi, A
Setiarto, A, Strüssmann, C A, Takashima, F, Watanabe, S & Yokota, M. Short-term responses of adult kuruma shrimp *Marsupenaeus japonicus* (Bate) to environmental salinity: osmotic regulation, oxygen consumption and ammonia excretion, 669
Shao, M Y *see* Kang, K H
Shelby, R A, Schrader, K K, Tucker, A, Klesius, P H & Myers, L J. Detection of catfish off-flavour compounds by trained dogs, 888
Shelby, R A, Shoemaker, C A & Klesius, P H. Development of an ELISA to measure the humoral immune response of hybrid striped bass *Morone chrysops* × *M. saxatilis* to *Streptococcus iniae*, 997
Shin, I-S *see* Choi, S-M
Shinn, A P *see* Dixon, B A
Shiozaki, K, Nakano, T, Yamaguchi, T, Sato, M & Sato, N. The protective effect of stevia extract on the gastric mucosa of rainbow trout *Oncorhynchus mykiss* (Walbaum) fed dietary histamine, 1421
Shirdhankar, M M, Thomas, P C & Barve, S K. Phenotypic estimates and heritability values of *Artemia franciscana*, 35
Shoemaker, C A *see* Shelby, R A
Sigholt, T *see* Ng, W-K
Silverstein, J T, Rexroad III, C E & King, T L. Genetic variation measured by microsatellites among three strains of domesticated rainbow trout (*Oncorhynchus mykiss*, Walbaum), 40
Simpson, R *see* Liu, W
Skapura, D P *see* Tiersch, T R
Sklan, D, Prag, T & Lupatsch, I. Structure and function of the small intestine of the tilapia *Oreochromis niloticus* × *Oreochromis aureus* (Teleostei, Cichlidae), 350
Sklan, D, Prag, T & Lupatsch, I. Apparent digestibility coefficients of feed ingredients and their prediction in diets for tilapia *Oreochromis niloticus* × *Oreochromis aureus* (Teleostei, Cichlidae), 358
Smith, T K *see* Tapia-Salazar, M
Sommerville, C *see* Dixon, B A
Southgate, P C *see* Acosta-Salmón, H
Sridhar, S *see* Haniffa, M A
Strüssmann, C A *see* Setiarto, A
Sullivan, C V, *see* Luckenbach, J A
Surendran, P K *see* Lalitha, K V
Szumiec, M A. Thermal limitation of carp (*Cyprinus carpio* L.) larvae survival in ponds of the temperate climate, 601

Takashima, F *see* Setiarto, A
Taller, J *see* Müller, T
Tan, B *see* Zheng, D
Tanaka, M *see* Islam, Md. S
Tapia-Salazar, M, Smith, T K, Harris, A, Cruz-Suárez, L-E & Ricque-Marie, D. Response of blue shrimp *Litopenaeus stylirostris* (Pérez-Farfante & Kensley, 1997) to dietary cadaverine supplementation, 1092
Tauber, R *see* Burridge, L E
Tendencia, E A. The first report of *Vibrio harveyi* infection in the sea horse *Hippocampus kuda* Bleekers 1852 in the Philippines, 1291
Teodorowicz, M *see* Krzemieniewski, M
Teshima, S-i *see* Alam, M S
Teugels, G G *see* Nguenga, D
Thomas, P C *see* Shirdhankar, M M
Thompson, K R, Muzinic, L A, Engler, L S, Morton, S-R & Webster, C D. Effects of feeding practical diets containing various protein levels on growth, survival, body composition, and processing traits of Australian red claw crayfish (*Cherax quadricarinatus*) and on pond water quality, 659
Thompson, P J *see* Jackson, C
Thórarinsdóttir, G G, *see* Jónasson, J P
Thunjai, T, Boyd, C E & Boonyaratpalin, M. Bottom soil quality in Tilapia ponds of different age in Thailand, 698
Tibbetts, S M, Lall, S P & Milley, J E. Apparent digestibility of common feed ingredients by juvenile haddock, *Melanogrammus aeglefinus* L., 643
Tičina, V, Grubišić, L & Katavić, I. Sampling and tagging of live bluefin tuna in growth-out floating cages, 307
Tidwell, J H *see* Coyle, S D
Tiersch, T R, Wayman, W R, Skapura, D P, Neidig, C L & Grier, H J. Transport and cryopreservation of sperm of the common snook, *Centropomus undecimalis* (Bloch), 278
Tinman, S, *see* Cnaani, A
Toro, J E, Alcapán, A C, Vergara, A M & Ojeda, J A. Heritability estimates of larval and spat shell height in the Chilean blue mussel (*Mytilus chilensis* Hupe 1854) produced under controlled laboratory conditions, 56
Toro, J E, Ojeda, J A & Vergara, A M. The genetic structure of *Mytilus chilensis* (Hupé 1854) populations along the Chilean coast based on RAPDs analysis, 1466
Trasviña, A *see* Peña, R
Treasurer, J W & Wadsworth, S L. Interspecific comparison of experimental and natural routes of *Lepeophtheirus salmonis* and *Caligus elongatus* challenge and consequences for distribution of chalimus on salmonids and therapeutant screening, 773
Troisi, G M *see* Rigos, G
Trzebiatowski, R *see* Rzemieniecki, A
Tsoumani, M *see* Paschos, I
Tucker, A *see* Shelby, R A
Turchini, G M, Mentasti, T, Crocco, C, Sala, T, Puzzi, C, Moretti, V M & Valfrè, F. Effects of the extensive culture system as finishing production strategy on biometric and chemical parameters in rainbow trout, 378
Tveit, K *see* Hemre, G-I
Tveiten, H, Jobling, M & Andreassen, I. Influence of egg lipids and fatty acids on egg viability, and their utilization during embryonic development of spotted wolf-fish, *Anarhichas minor* Olafsen, 152
Tyrpenou, A E *see* Rigos, G
Tzchori, I, Degani, G, Elisha, R, Eliyahu, R, Hurvitz, A, Vaya, J & Moav, B. The influence of phytoestrogens and oestradiol-17β on growth and sex determination in the European eel (*Anguilla anguilla*), 1213

Urbanyi, B *see* Lahnsteiner, F
Urbinati, E C *see* Martins da Rocha, R
Ureta, J P *see* Estay, F J

Vaccaro, A M, Buffa, G, Mirto, S, Sarà, G & Mazzola, A. Comparison of growth performance and biometric relationships in two reciprocal sturgeon hybrids reared in net cages (Sicily, Mediterranean), 552
Valfrè, F *see* Turchini, G M
van de Vis, H. Safety and Quality Issues in Fish Processing, 100
van der Meeren, T & Harboe, T. Live Feeds in Marine Aquaculture, 213
Vaya, J *see* Tzchori, I
Venkat, H K, Sahu, N P & Jain, K K. Effect of feeding Lactobacillus-based probiotics on the gut microflora, growth and survival of postlarvae of *Macrobrachium rosenbergii* (de Man), 501
Vera-Cartas, J *see* Schneider, O
Vergara, A M *see* Toro, J E
Véron, B *see* Jo, Q
Verreth, J A J *see* El-Shafai, S A
Verreth, J A J *see* Schneider, O
Vielma, J, Ruohonen, K, Gabaudan, J & Vogel, K. Top-spraying soybean meal-based diets with phytase improves protein and

© 2004 Blackwell Publishing Ltd

mineral digestibilities but not lysine utilization in rainbow trout, *Oncorhynchus mykiss* (Walbaum), 955
Vilchez, C *see* Lee, K-J
Villarreal, H, Hernandez-Llamas, A, Rivera, M C, Millan, A & Rocha, S, Effect of substitution of shrimp meal, fish meal and soy meal with red crab *Pleuroncodes planipes* (Stimpson) meal in pelleted diets for postlarvae and juvenile *Farfantepenaeus californiensis* (Holmes), 178
Villarreal-Colmenares, H *see* Cortés-Jacinto, E
Vita, R, Marín, A, Jiménez-Brinquis, B, Cesar, A, Marín-Guirao, L & Borredat, M, Aquaculture of Bluefin tuna in the Mediterranean: evaluation of organic particulate wastes, 1384
Vogel, K *see* Vielma, J
Voltolina, D *see* Lora-Vilchis, M C
Vromant, N, Nam, C Q, Chau, N T H & Ollevier, F, Survival rate and growth performance of *Cyprinus carpio* L. in intensively cultivated rice fields, 171
Vught, I *see* Harzevili, A S

Waddy, S L *see* Burridge, L E
Wadsworth, S L *see* Treasurer, J W
Wahab, M A *see* Alim, M A
Walden, J *see* Fitzsimmons, S
Wang, X *see* Choi, S-M
Wang, X *see* Kim, K-W
Wang, X *see* Li, P
Wang, Z *see* Zhuang, S
Warner-Smith, R J *see* Booth, M A
Watanabe, S *see* Setiarto, A
Wayman, W R *see* Tiersch, T R
Webster, C D *see* Coyle, S D
Webster, C D *see* Thompson, K R
Welker, T L & Congleton, J L, Oxidative stress in juvenile chinook salmon, *Oncorhynchus tshawytscha* (Walbaum), 881
Wen, H S & Lin, H R, Effects of exogenous neurohormone, gonadotropin (GtH) and dopaminergic drugs on the serum GtH content and ovulatory responsiveness of wild catfish, *Silurus asotus* (Linnaeus, 1758), 204
Westcott, J D, Hammell, K L & Burka, J F, Sea lice treatments, management practices and sea lice sampling methods on Atlantic salmon farms in the Bay of Fundy, New Brunswick, Canada, 784
Whoriskey, F *see* Carr, J
Willis, K J & Ling, N, Toxicity of the aquaculture pesticide cypermethrin to planktonic marine copepods, 263
Winkler, F M, García-Melys, D & Palma-Rojas, C, Karyotypes of three South East Pacific flounder species of the family Paralichthyidae, 1294
Wislocki, P G *see* Burridge, L E
Wissing, P J *see* Gensic, M
Wong, W H & Levinton, J S, Culture of the blue mussel *Mytilus edulis* (Linnaeus, 1758) fed both phytoplankton and zooplankton: a microcosm experiment, 965

Xie, S *see* Han, D
Xie, S *see* Xue, M
Xu, W *see* Zheng, D
Xue, M, Xie, S & Cui, Y, Effect of a feeding stimulant on feeding adaptation of gibel carp *Carassius auratus gibelio* (Bloch), fed diets with replacement of fish meal by meat and bone meal, 473

Yamaguchi, T, *see* Shiozaki, K
Yamamoto, S *see* Hattori, M
Yang, H, & Guo, X, Tetraploid induction by meiosis inhibition in the dwarf surfclam *Mulinia lateralis* (Say 1822): effects of cytochalasin B duration, 1187
Yang, Y *see* Han, D
Yang, Y, *see* Lin, H Z
Yildiz, H Y *see* Benli, A C K
Yokota, M *see* Setiarto, A

Zamora, A G *see* Patiño-Suárez, V
Zamorano, M J *see* Ginés, R
Zbanyszek, R *see* Halver, J E
Zhang, M *see* Zhuang, S
Zhang, W *see* Zheng, D
Zhang, X *see* Zhuang, S
Zhang, Z F *see* Kang, K H
Zheng, D, Mai, K, Liu, S, Cao, L, Liufu, Z, Xu, W, Tan, B & Zhang, W, Effect of temperature and salinity on virulence of *Edwardsiella tarda* to Japanese flounder, *Paralichthys olivaceus* (Temminck et Schlegel), 494
Zheng, W, *see* Lin, H Z
Zhu, X *see* Han, D
Zhuang, S, Zhang, M, Zhang, X & Wang, Z, The influence of body size, habitat and diet concentration on feeding of *Laternula marilina* Reeve, 622
Zolota, V *see* Poulos, C


© 2004 Blackwell Publishing Ltd

# Keyword index


abalone, 1062, 1076
absorption, 350
acetylcholinesterase activity, 1310
*Acipenser*
  *persicus*, 1238
  *ruthenus*, 144, 519
acrosome, 519
acute toxicity, 1106
adaptation, 473
adhesion kinetics, 403
AFLPs, 1472
African catgish, 232
agroecology, 289
allozyme electrophoresis, 1328
alternate ingredients, 178
alternative protein ingredients, 1
amino acids, 312, 1110
ammonia, 134
ammonia excretion, 669
amoebic gill disease, 112, 1448
amylase, 1158
anaesthesia, 1025
Anarhichadidae, 152
*Anarhichas lupus*, 613
animal
  breeding, 924
  by-products, 1
  protein sources, 1261
apparent digestibility coefficients, 1, 1261, 1441
aquaculture, 465, 483, 552, 751, 1117, 1187
aquaculture environment, 629
aquatic epidemiology, 764
arginine, 842
*Argopecten irradians*, 678
*Artemia*, 35, 62, 547, 607, 1086
artemia, 501
artifical
  spawning, 278
  insemination, 1486
  spawning, 1318
  substrate, 817
ascorbic acid, 256
assimilation efficiency, 622
astaxanthin, 458
Atlantic
  cod, 453, 683, 992
  salmon, 112, 1228, 1448
  salmon *Salmo salar L.*, 784

*Bacillus* sp., 1441
band-sharing, 807
*Barbodes gonionotus*, 171
barley-based fermented grains, 1158
battery culture, 559
bighead carp *Aristichthys nobilis* (Richardson), 836
bioavailability, 1299
biochemical composition, 1131
bioeconomics, 103
bioencapsulation, 943
biogenic capacity, 616
biometric relationship, 552
biopsy, 96
black sea bass, 1457
blood
  glucose, 874
  parameters, 859, 1390
  plasma, 1025
Bluefin tuna, 1387
bluefin tuna (*Thunnus thynnus* L.), 310
body
  composition, 1207
  movements, 453
  shape, 924
  weight, 893
bottom soils, 698
*Brachionus plicatilis*, 1394
*Brachionus*, 62
brackish water earthen pond, 423
brewer's by-products, 1
British Columbia, 483
broodstock, 99
burbot, 49

cadmium, 80
*Caligus elongatus*, 764, 773, 784
Cameroon, 1349, 1358
canine olfaction, 888
canola, 15, 25, 981
capture, 245
carbohydrates, 1220
carbon, 194
caridean shrimp, 1179
carotenoids, 905
carp, 1318
carp larvae, 604
catfish, 99, 204, 888
catfish diets, 917
catheter, 87
*Catla catla*, 874
catla, 124
CB, 1062
*Centropages hamatus*, 432
centrum defects, 850
*Chaetoceros muelleri*, 1086
chalimus, 773
*Charonia* sp., 1202
chemotherapeutants, 773
*Cherax*, 905
*Cherax*
  *destructor*, 1328
  *destructor* Clark, 1106
  *quadricarinatus*, 71, 659
  *quadricarinatus* (von Martens), 559
Chile, 56
Chinese longsnout catfish, 866
chinook salmon, 881


© 2004 Blackwell Publishing Ltd

*Chlamys*
*farreri*, 678
*islandica*, 1405
chloramine-T, 112, 1448
chlorine dioxide, 112
cholesterol, 529
chymotrypsin, 914
ciliate, 518
*Cirrhinus*
*mrigala*, 162
*mrigala* H., 134
*Clarias gariepinus*, 232
Clariidae, 1349, 1358
clearance rate, 622
clonal lines, 1472
Close up Solidus P, 250
*Clostridium botulinum*, 594
clove oil, 1025
clown fish, 1009
cold tolerance, 893
commercial, 1039
common
carp, 124
dentex (*Dentex dentex*), 224
constant magnetic field (CMF), 568
copepod parasite, 730
copepods, 263, 432, 742
correlations, 529
cortisol, 245
cottonseed meal, 859
*Crassostrea gigas*, 1131
crayfish, 559
crowding, 245
crustacean, 1166, 1172
cryopreservation, 519, 1195, 1415, 1429, 1457
*Ctenopharyngodon idella*, 1277
cuttlefish, 1141
cypermethrin, 263
*Cyprinus carpio*, 171, 547
cytochalasin B, 1187

deep-water chlorophyll maxima, 1253
defatted soybean meal, 299
dehulled soybean meal, 410
delay, 607
delayed growth, 1141
desiccation, 759
DHA, 1394
diapause eggs, 432
dietary, 80
dietary
cadaverine, 1092
lipids, 709
protein, 1277
requirement, 842
threonine requirement growth studies, 162
diets, 436, 659, 683, 965, 1039, 1179
digestibility, 312, 358, 643, 981, 1158, 1228, 1370
digestible energy, 970
digestion, 350, 1149
digestive
enzymes, 613
proteases, 914
*Diplodus puntazzo*, 1220
diploidization, 271

discharge load, 1053
disinfection, 112
distillers grains, 365
distribution, 793
diversification, 289
6-DMAP, 1076
DNA, 807
dry duckweed, 574
duckweed digestibility, 574
duration effect, 432

E ratio, 250
early life stages, 263
EDTA, 1339
*Edwardsiella tarda*, 494, 1390
effluent water, 321
eggs, 224, 759, 1277
egg
incubation temperature, 1358
quality, 152
storage temperature, 1358
surface disinfectants, 992
ELISA, 997
emamectin benzoate, 713, 784
embryonic development, 224
energy, 250
energy budget, 866
enrichment, 1394
enzyme, 134
*Epinephelus coioides*, 1261
epiphytes, 508
eqq quality, 613
erythrocyte, 874
estradiol-17β, 397, 1213
ethylene glycol, 1415
eugenol, 1005, 1025
European
eel, 1213
lobster, 607
sea bass, 636
sheatfish *Silurus glanis* L., 568
exotoxin, 1395
extensive farming, 378
extracellular products, 931, 1395
extrusion, 358
extrusion processing, 1153
eye structure, 345

faeces recovery, 1370
*Farfantepenaeus californiensis*, 178
farmed freshwater prawn, 629
farming, 310
fatty acids, 547, 1039, 1131, 1220, 1228
FCM, 1062, 1076
fecundity, 1031
feed
conversion efficiency, 1305
efficiency, 1305
efficiency ratio, 1141
intake, 1305
utilization, 859
feeding, 607, 622, 965
feeding
rate, 1141
stimulant, 473


© 2004 Blackwell Publishing Ltd

feminization, 397
filefish, 1429
fillet
composition, 709
quality, 378, 1039
fingerling, 604
finishing, 378
first
feeding, 345
intestinal bacteria, 436
fish, 144, 403, 1305
fish
culture, 289
farming, 764
farms, 793
meal replacement, 828
meal, 410, 473, 859, 1092, 1153
nutrition, 709
oils, 709, 1394
rearing, 568
fishmeal replacement, 15, 25
fishmeal, 299, 1370
floating cage, 310
flocs, 483
flounder, 250
flounder karyotypes, 1298
food
conversion efficiency, 1039
value, 1086
web, 124
formulated diets, 1244
fresh duckweed, 574
freshwater crayfish, 1328

gadoids, 643
gastric mucosa, 1421
Gastropod, 1149
genetic
diversity, 587
resources, 278
structure, 1466
variation, 1305
genistein, 1213
genotype-by-environment interaction, 924
geosmin, 888
germplasm, 278
geutathione redox system, 256
gibel carp, 473
gilthead sea bream, 828
glucose, 245
glutathione peroxidase, 1099
GnRHa implant, 1486
gonad, 96
gonadotropin, 204
*Gracilariopsis bailinae*, 423
grouper, 1261
growth, 80, 99, 215, 436, 501, 559, 616, 636, 652, 659, 1013, 1039, 1131, 1158, 1207, 1270
growth
compensation, 836
performance, 552, 828
rate, 423, 1086
response, 859
variation, 559
Gynogenesis, 271, 1472

haddock, 87, 465, 643, 992
haematocrit, 529
haematological parameters, 80
haematology, 134, 1310
haemoglobin, 529, 874
halibut larvae, 1286
*Haliotis rubra*, 1062, 1076
haloperidol, 10
hatching, 432
hatching time, 453
hCG, 144
heat shock, 937
hepatopancreas, 1172
heritability, 35, 56
*Heterobranchus longifilis*, 1349, 1358
*Heteropneustes fossilis*, 937
heterotrophic culture, 371
*Hippoglossina macrops*, 1298
histamine, 1421
histopathology, 931
*Homarus americanus*, 713
hormonal sex reversal, 397
hormones, 1025
HPI-axis, 1117
HUFAs, 152
husbandry, 1244
hybrid, 916
hybrid striped bass, 997, 1383
hybridization, 99
hydrilla, 1277
hydrogen peroxide, 112

Iceland scallop, 1405
ideal protein, 1110
immune parameter, 678
immunoglobulin, 997
immunostimulant, 1383
inbreeding, 340
indicator bacteria, 629
induced spawning, 1486
induction of ovulation, 10
ingestion, 62, 1149
ingestion rate, 518, 622
inheritance, 1328
innate immunity, 1434
inoculation, 436
integrated agriculture–aquaculture systems, 289
intensive culture, 1039
iodine, 759, 1117
ionic parameters, 1238
iron, 256, 859
*Isochrysis* sp., 1086

juvenile quality, 613

koi carp, 10
kuruma shrimp, 669, 842

*L. acidophillus*, 501
*L. sporogenes*, 501

*Labeo rohita* fingerlings, 436
largemouth bass, 312
larva, 1149

© 2004 Blackwell Publishing Ltd 

larvae, 518
larval
development, 1179
feeding, 547
growth, 56
*Laternula marilina*, 622
$LC_{50}$, 713, 1106
leaching, 842
Lecirelin, 1318
*Leiocassis longirostris* (Günther), 866
*Lepeophtheirus*
*salmonis*, 730, 759, 773, 784
*salmonis* (Krøyer), 742
*Lepidium meyenii*, 215
*Lepomis macrochirus*, 397
*Leucaena leucocephala*, 436
leukocyte, 874
levamisole, 1383
LHRH, 144
LHRH-A, 204
LHRHa, 10
licorice, 1213
light, 49
light
intensity, 345
regime, 683
liming, 698
linkage map, 893
lipid
oxidation, 948
peroxidation, 881
sources, 1228
*Listonella anguillarum*, 403
*Litopenaeus*
*stylirostris*, 103, 1244
*vannamei*, 321, 518, 1153, 1158, 1441
lobster, 713
lupin seed meal, 828
lupins, 981
lysine digestibility, 955
*Lysmata* sp., 1179

maca, 215
*Macrobrachium rosenbergii*, 501, 629, 943, 1415
mahseer, 1270
malondialdehyde, 256
mariculture, 184
marine, 1339
marine
animal protein, 1158
by-products, 643, 1153
fish, 87
ornamental aquaculture, 1179
*Marsupenaeus japonicus*, 669, 842
matrinxã, 245
maturation, 905
MDA, 1310
mean
corpuscular haemoglobin concentration, 529
corpuscular haemoglobin, 529
corpuscular volume, 529
meat and bone meal, 473
meiosis, 1187
mekong delta, 1013
*Melanogrammus aeglefinus*, 465
metabolic enzymes, 613
metals, 652
metamorphosis, 616
2-methylisoborneol, 888
17α-methyl testosterone, 943
microalgae, 1131
microbiology, 184
microflora, 629
*Micropterus salmoides*, 312
microsatellite marker variation, 40
microsatellites, 1472
milkfish, 423
mineral, 955
mola, 124
molecular taxonomy, 1009
mollusk, 678
monoclonal antibody, 997
monosex, 71
*Morone*, 997, 1383
morphology, 99, 807
mortality, 1405
motility, 519
MS-222, 1025
mtDNA, 1328
mud crab, 1013
*Mulinia lateralis*, 1187
mullet, 817
Murray cod *Maccullochella peelii peelii*, 1039
muscle quality parameters, 683
mussel, 965
mussel growth, 1253
*Mytilus chilensis*, 56, 1466
*Mytilus edulis*, 965

natural
biota, 194
feed, 194
*Neoparamoeba pemaquidensis*, 1448
Nile tilapia, 271, 1110, 1390, 1472
nitrogen, 194, 817, 1053
Northeast Thailand, 289
nutrient balance, 1370
nutrients, 483, 1053
nutrition, 178, 194, 970, 1149, 1166, 1172

off-flavour, 888
olive flounder, 410
*Oncorhynchus*
*mykiss*, 40, 378
*tshawytscha*, 1099
ontogeny, 152
*Oreochromis*, 340, 1434
*Oreochromis*
*aureus*, 1031
*niloticus*, 171, 271, 574, 587, 1110, 1370, 1390, 1472
*spilurus*, 1031
osmolality, 1238, 1310
osmotic regulation, 669
ovulation, 204
oxidative stress, 881
oxolinic acid, 1299
oxygen
consumption, 669, 1005
deficiency, 850


© 2004 Blackwell Publishing Ltd

P fractionation, 1339
*P. adspersus*, 1298
*P. microps*, 1298
*Pagrus*
*auratus*, 458
*major*, 850
*Panagrellus redivivus*, 547
*Paralabrax maculatofasciatus*, 345
*Paralichthys*
*lethostigma*, 1486
*olivaceus*, 494, 1310
parent–offspring regression, 35
particulate waste, 1387
pathogenesis, 1395
PCR, 807
pearl oyster, 385
peas, 981
*Pelodiscus sinensis*, 948
penaeid shrimp, 1092
peppermint shrimp, 1179
pepsin, 1421
*Perca fluviatilis*, 1005
performance, 1110
*Perna canaliculus*, 1253
pharmacokinetics, 1299
phenanthrene, 1310
phenotypic estimates, 35
phosphate, 1339
phosphorus, 312, 1053
phosphorus requirement, 636
*Photobacterium damsela* subsp. *piscicida*, 931
photoperiod, 465, 1207
phytase, 955
phytooestrogens, 1213
pikeperch, 916
*Pinctada*, 96
*Pinctada imbricata*, 385
pituitary homogenate, 1318
plant
protein sources, 1261
proteins, 15, 25, 643
*Pleuronectes planipes*, 178
polyculture, 124, 171, 423
*Pomoxis nigromaculatus*, 397
ponds, 1339
population differentiation, 40
prawn, 1415
predation, 1202
prey, 1202
probiotics, 501
production, 71, 1013
production
traits, 256
rate, 423
protein, 71, 250, 365, 659, 955, 970
protein
efficiency, 970
replacement, 917
requirement, 1270
sources, 312
proximate composition, 1277
pulse exposures, 263
punti, 124

QTL, 893

quality, 184
quantitative genetics, 924
quarantine, 1244

raceway culture, 529
radioimmunoassay, 943
rainbow trout, 40, 215, 378, 1421
rainbow trout
of the Kamloops strain, 529
*Oncorhynchus mykiss*, 1
*Rana catesbeiana*, 616
RAPD, 916, 1009
RAPD (DNA), 587
RAPDs, 730, 1328, 1466
rapeseed, 15, 25
ration level, 866
raw material freshness, 1092
recruitment, 385
red
blood cell count, 529
claw, 659
colour, 340
sea bream, 850
tilapia, 1031
reference values, 529
regulation, 321
replacement, 410
reproduction, 144, 152, 385, 465, 1166, 1486
reproductive performance, 836, 1277, 1349
residual feed intake, 1305
retinoids, 905
rice–fish culture, 171
risk assessment, 103
rohu, 124
roquette seed meal, 917
rotifer, 1394
*Rutilus rutilus*, 1005

*S. gigas*, 1149
salinity, 494, 669, 914, 1286
*Salmo salar*, 730, 773, , 759
salmon, 594, 764
salmon
farming, 263, 742
lice, 751
salmonids, 709, 793, 924, 1005
sampling and tagging mortality, 310
sander
lucioperca, 916
volgense, 916
*Scylla*, 1013
*Scylla serrata* larvae, 62
sea
bass, 184, 931
bream, 184
cages, 1448
horse, 1294
lice, 723, 730, 742, 784
lice larvae, 793
seabream, 15, 25
season, 683
*Sebastes schlegeli*, 80
Sedex, 1339
sediment
sediments, 184, 652

© 2004 Blackwell Publishing Ltd

seed production, 1031
selection, 340
selective breeding, 40
selenium, 1099
semen, 232
seminal fluid, 1238
serum protein, 874
sex
  determination, 1213
  differentiation, 1213
  ratio, 1328
  reversal, 943
SGR, 501
shade, 458
sharpsnout sea
  bream, 1220
  bream *Diplodus puntazzo*, 1299
short-term storage, 232, 1195
shrimp, 194, 1244, 1395
shrimp
  effluent, 817
  farm, 321, 1053
  larvae, 652
siganids, 817
*Silurus*, 99
silver perch, 970, 981
size rank, 559
skin colour, 458, 924, 1207
SLICE®, 784
SLICE™, 713
small intestine, 350
smolts, 1099
snapper, 15, 25, 299
social interaction, 559
sodium hypochlorite, 759
soft-shelled turtle, 948
soil carbon, 698
somitogenesis, 850
soybean, 1220
soybean meal, 917, 955, 981
*Sparus aurata*, 914, 1207
spat, 1131
spawning, 1031
species comparison, 1434
sperm, 144, 1195, 1415, 1429, 1457
sperm motility, 1238
sperm-to-egg ratio, 1486
spermatophore, 1415
spermatozoa, 519
stable isotopes, 508
steelhead trout, 1117
sterlet, 519
steroid, 1457
sterolic composition, 371
stevia extract, 1421
strain comparison, 1349, 1358
*Streptococcus iniae*, 997
stress, 245, 1099, 1117, 1286
stress response, 881, 1434
stunting, 836
sturgeon, 144, 552
substrate, 616
surface components, 403
survey, 784
survival, 215, 501, 604, 652, 659, 1286
survival analysis, 1405
suspended sediment, 483

temperature, 49, 224, 453, 494, 604, 678, 1131, 1405
temperature effect, 432
tetraploidy, 937, 1187
*Tetraselmis suecica*, 371
*Thamnaconus septentrionalis*, 1429
Tilapia, 340, 350, 358, 365, 698, 893, 1370, 1434
time-series analysis, 764
α-tocopherol, 1421
α-tocopheryl acetate, 948
tolerance, 1405
*Tor putitora*, 1270
total
  plasma protein, 529
  suspended solids, 1053
toxicity, 112, 134, 713
toxins, 594
trace elements, 483
transportation, 1005
tri-acyl-*sn*-glycerols, 1394
trichlorfon, 1106
triploidy, 937, 1062, 1076, 1187
tropical highlands climate, 1349
trout, 1025
trumpet shell, 1202
trypsin, 914
tuna fattening, 1387

urechidae, 1195
*Urechis unicinctus*, 1195
urinary system, 87
urine collection, 87

vegetable oils, 709
*Vibrio harveyi*, 1294
*Vibrio*, 1395
Vietnam, 1013
virulence, 494
virulence products, 1395
vitamin E, 948, 1172

water, 184
water
  quality, 568, 659
  temperature, 1228
  treatment, 568
weather, 604
weight gain, 1305
wheat gluten meal, 1158
wild
  and escaped farmed and landlocked salmon, 723
  salmonids, 751
Wolffish, 613

yabby, 1106, 1328

zooplankton, 1106



© 2004 Blackwell Publishing Ltd